

## Anaesthesia recommendations for patients suffering from

### **Mounier-Kuhn syndrome**

**Disease name:** Mounier-Kuhn syndrome

**ICD 10:** Mounier-Kuhn syndrome Q32.4; Q32.1

- With bronchiectasis: J47
- Exacerbation (acute): J47
- Lower respiratory infection: J47
- Acquired: J98.09
- With bronchiectasis: J47
- With exacerbation (acute): J47
- With lower respiratory infection: J47.0

**Synonyms:** Tracheobronchomegaly, trachiectasis, tracheobronchomalacia, and multiple tracheal diverticula.

A marked dilation of the trachea and main bronchi, associated with a higher rate of recurrent lower respiratory tract infections, characterizes the Mounier-Kuhn syndrome (MKS). The clinical presentation varies widely from minimal respiratory discomfort, repetitive respiratory infections, mucus accumulation and ineffective cough, to severe respiratory failure. This syndrome is considered to be congenital, although most of the times it presents in the third or later decades with recurrent respiratory tract infections. The aetiology is uncertain, anatomic-pathological findings have led to believe it due to the lack of smooth muscle and elastic connective tissue in the trachea and main bronchi, causing tracheobronchomegaly and "herniation" of diverticula between the cartilaginous rings.

---

Medicine in progress



Perhaps new knowledge

Every patient is unique

Perhaps the diagnostic is wrong

---



Find more information on the disease, its centres of reference and patient organisations on Orphanet: [www.orpha.net](http://www.orpha.net)

---

### **Disease summary**

---

Frequently the diagnosis can be sensed based on in case of abnormal chest X-ray, and an anaesthesiologist should be suspected it when there is an unexplainable loss of air during ventilation with tracheal tube, but it needs to be confirmed by a CT scan to allow for precise measurement of airways and evaluation of additional changes in pulmonary tissues and possibly a bronchoscopy could be considered for additional evaluation.

The treatment does not include surgery and it is typically medical includes supportive measures and possible airway stenting in cases of significant tracheomalacia, but surgery has also been attempted. The only clinical trial which included 12 patients with Mounier-Kuhn syndrome showed significant improvements in pulmonary function and quality of life when compared to baseline. The objective of conservative treatment is sputum clearance, using positional physiotherapy and early and aggressive treatment of pulmonary infections. In some cases chronic prophylactic administration of antibiotics may be required.

The diagnosis can be sensed in case of abnormal chest X-ray, and should be suspected when there is an unexplainable loss of air during ventilation with tracheal tube, but needs to be confirmed by a CT scan and possibly a bronchoscopy.

---

### **Typical surgery**

---

Surgery is not generally indicated, but in some cases of severe tracheomalacia and expiratory collapse of trachea a tracheobronchoplasty has been used, and in refractory end stage lung disorder lung transplant is the definitive treatment.

---

### **Type of anaesthesia**

---

According to the type of surgery, lack of the “seal” in case of tracheal intubation should be concealed.

---

### **Necessary additional diagnostic procedures (preoperative)**

---

CT scan, bronchoscopy.

---

### **Particular preparation for airway management**

---

We suggest a thorough study of the CT scan and the measurement of the tracheal diameters, especially in the sections where the cuff of the endotracheal tube will be placed, to choose the size and the most suitable position for the tracheal tube.

Before induction, a set of different sized tubes, several supraglottic devices, a fiberscope, and a manometer to measure the pressure of the cuff should be prepared.

After endotracheal intubation, a small ET tube could be inserted in the oesophagus, cuffed and a stomach aspiration tube should be inserted into it to oesophageal tube, in order to reduce the risk of aspiration.

---

#### **Particular preparation for transfusion or administration of blood products**

---

According to type of surgery.

---

#### **Particular preparation for anticoagulation**

---

According to type of surgery.

---

#### **Particular precautions for positioning, transport or mobilisation**

---

Not indicated.

---

#### **Probable interaction between anaesthetic agents and patient's long term medication**

---

Not reported.

---

#### **Anaesthesiologic procedure**

---

Before induction, a set of different sized tubes, several supraglottic devices, a fiberscope, and a manometer to measure the pressure of the cuff should be prepared.

A rapid sequence intubation should be performed in order to avoid gastric distension.

After endotracheal intubation, a small ET tube could be inserted in the oesophagus, cuffed and a stomach aspiration tube should be inserted into it to oesophageal tube, in order to reduce the risk of aspiration.

Should the ventilator detect an air loss, it is possible to place an extra-gloctic device (I-gel, Fastrach) and a small ET tube through it, with the purpose of enhancing the seal with the esophageal cuff.

---

#### **Particular or additional monitoring**

---

Pay particular attention to the pressure of the ET cuff to avoid mechanical stress on the tracheal wall.

Monitor strictly the Pressure/Volume loop on the ventilator in order to detect any air loss.

---

### **Possible complications**

---

Air loss during ventilation.

Tracheal lesions or even rupture in case of excessive pressure from the cuff or during intubation.

Aspiration pneumonia.

---

### **Postoperative care**

---

Strict chest X-ray monitor to detect early pulmonary complication from inhalation or atelectasis.

Early mobilisation and positional physiotherapy. Prophylactic antibiotic therapy in case of respiratory infections.

---

### **Information about emergency-like situations / Differential diagnostics**

---

*caused by the illness to give a tool to distinguish between a side effect of the anaesthetic procedure and a manifestation of the disease*

None.

---

### **Ambulatory anaesthesia**

---

See post-operative care.

---

### **Obstetrical anaesthesia**

---

If possible avoid general anaesthesia.

## Literature and internet links

1. Cataldo R, Galli B, Proscia P, Carassiti M. Management of a patient with Mounier-Kuhn syndrome undergoing repeated general anesthetics. *Can J Anaesth.* 2013 Jun;60(6):602-3. doi: 10.1007/s12630-013-9917-7
2. Weissleder R, Wittenberg J, Harisinghani MG et-al. *Primer of diagnostic imaging.* Mosby Inc. (2007) ISBN:0323040683
3. Mcguinness G, Naidich DP, Leitman BS et-al. Bronchiectasis: CT evaluation. *AJR Am J Roentgenol.* 1993;160 (2): 253-9
4. Cartier Y, Kavanagh PV, Johkoh T et-al. Bronchiectasis: accuracy of high-resolution CT in the differentiation of specific diseases. *AJR Am J Roentgenol.* 1999;173 (1):47-52
5. Barker AF. Bronchiectasis. *N. Engl. J. Med.* 2002;346 (18):1383-93
6. Cantin L, Bankier AA, Eisenberg RL. Bronchiectasis. *AJR Am J Roentgenol.* 2009;193 (3): W158-71. doi:10.2214/AJR.09.3053
7. Solanki T, Neville E. Bronchiectasis and rheumatoid disease: is there an association? *Br. J. Rheumatol.* 1992;31(10):691-3
8. Fenlon HM, Doran M, Sant SM et-al. High-resolution chest CT in systemic lupus erythematosus. *AJR Am J Roentgenol.* 1996;166 (2):301-7
9. Collins J, Stern EJ. *Chest radiology, the essentials.* Lippincott Williams & Wilkins. 2007; ISBN:0781763142
10. Naidich DP, Srichai MB, Krinsky GA. *Computed tomography and magnetic resonance of the thorax.* Lippincott Williams & Wilkins. 2007; ISBN:0781757657
11. Warrell DA. *Oxford textbook of medicine, Sections 18-33.* Oxford University Press, USA. 2005; ISBN:0198569785
12. Javidan-Nejad C, Bhalla S. Bronchiectasis. *Thorac Surg Clin.* 2010;20 (1):85-102. doi:10.1016/j.thorsurg.2009.12.007
13. Javidan-Nejad C, Bhalla S. Bronchiectasis. *Radiol. Clin. North Am.* 2009;47 (2):289-306. doi:10.1016/j.rcl.2008.11.006
14. Ooi GC, Khong PL, Chan-Yeung M, et al. High-resolution CT quantification of bronchiectasis: clinical and functional correlation. *Radiology.* 2002;225 (3):663-72. [Radiology \(full text\)](#) - doi:10.1148/radiol.2253011575
15. Mounier-Kuhn P. Dilatation de la trachée: constatations, radiographiques et bronchoscopies. *Lyon Med* 1932; 150:106–9
16. Spencer H. Congenital abnormalities of the lung; congenital tracheobronchomegaly. In: Spencer H, ed. *Pathology of the Lung*, 4th edition. Pergamon Press: Oxford, 129–30
17. Aaby GV, Blake HA. Tracheobronchomegaly. *Ann ThoracSurg* 1966;2:64–70
18. Himalstein MR, Gallagher JC. Tracheobronchomegaly. *Ann Otol Rhinol Laryngol* 1973;82:223–7
19. Wolfgang Dahnert ED. *Radiology Review Manual-second Edition,* William and Wilkins, 1993. Lippincot Williams and Wilkins: NY, USA
20. Bateson EM, Woo-Ming M. Tracheobronchomegaly. *ClinRadiol* 1973;24:354–8
21. Sorenson SM, Moradzadeh E, Bakhda R. Repeated infections in a 68-year-old man. *Chest* 2002; 121:644–6
22. Giannoni S, Benassai C, Allori O, et al. Tracheomalacia associated with Mounier-Kuhn syndrome in the intensive care unit: treatment with Freitag stent. A case report. *Minerva Anestesiol* 2004; 70: 651–9
23. Van Schoor J, Joos G, Pauwels R. Tracheobronchomegaly—the Mounier-Kuhn syndrome: report of two cases and review of the literature. *Eur Respir J* 1991;4:1303–6
24. Pilavaki M, Anastasiadou K, Vlachojanni E, et al. Tracheobronchomegaly (Mounier-Kuhn syndrome): roentgen findings and tracheal stent instrumentation. *Pneumologie* 1995;49:556–8
25. Odell D, Shah A, Gangadharan S, et al. Airway Stenting and Tracheobronchoplasty Improve Respiratory Symptoms in Mounier-Kuhn Syndrome. *Chest* 2011; 140(4):867–873. doi: 10.1378/chest.10-2010.

---

**Last date of modification: May 2014**

---

*This guideline has been prepared by:*

**Author**

**Rita Cataldo**, Department of Anaesthesia, Campus BioMedico, University School of Medicine, Rome, Italy  
[r.cataldo@unicampus.it](mailto:r.cataldo@unicampus.it)

**Peer revision 1**

**Sudhakar Subramani**, Department of Anaesthesia, University of Iowa, USA  
[sudhakar-subramani@uiowa.edu](mailto:sudhakar-subramani@uiowa.edu)

**Peer revision 2**

**Eduard Krustins**, Department of Internal Medicine, Pauls Stradins Clinical University Hospital, Riga, Latvia  
[eduards.krustins@stradini.lv](mailto:eduards.krustins@stradini.lv)

---